

SCHEDULING STATUS **S3**

1. NAME OF THE MEDICINE

MAINTENANCE SOLUTION FRESenius

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Each 1 000 ml contains:	
Potassium chloride	1,938 g
Sodium chloride	1,5198 g
Dextrose anhydrous	50,0 g or
Dextrose monohydrate	55,0 g

Content of electrolytes per 1 000 ml:	
sodium ion (Na ⁺)	26 mmol/l
potassium ion (K ⁺)	26 mmol/l
chloride ion (Cl ⁻)	52 mmol/l

Excipient with known effect:
Contains sugar (dextrose).
For the full list of excipients, see section 6.1.

3. PHARMACEUTICAL FORM

Solution for intravenous infusion.
Clear, colourless to light straw-coloured solution.

Osmolarity: 382 mOsm/l
pH: 4,5
Energy: 840 kilojoules.

4. CLINICAL PARTICULARS

4.1 Therapeutic indications

MAINTENANCE SOLUTION FRESenius is indicated for use in adults and paediatric patients as a source of electrolytes, energy and water.

4.2 Posology and method of administration

MAINTENANCE SOLUTION FRESenius is for intravenous use only.
The dosage is to be directed by a medical doctor and is dependent upon age, body mass, clinical condition of the patient and laboratory determinations. Frequent laboratory determinations and clinical evaluations are essential to monitor changes in blood glucose and electrolyte concentrations, and fluid and electrolyte balance during prolonged parenteral therapy.
When MAINTENANCE SOLUTION FRESenius is to be administered peripherally, it should be slowly infused through a small bore catheter, placed well within the lumen of a large vein to minimise venous irritation. Carefully avoid tissue infiltration.
Usually up to 40 mmol of potassium per litre daily is sufficient to replace normal loss in adults. Infusion rates should not exceed 10 mmol per hour or 120 mmol per day. Paediatric patients may require 2 to 3 mmol per kg of body mass daily.
Fluid administration should be based on calculated maintenance or replacement fluid requirements for each patient.
Some additives may be incompatible. A pharmacist should be consulted. When introducing additives, use aseptic techniques. Mix thoroughly. Do not store.
MAINTENANCE SOLUTION FRESenius should be inspected visually for particulate matter and discolouration prior to administration, whenever solution and container permit.

4.3 Contraindications

MAINTENANCE SOLUTION FRESenius is contraindicated where the administration of sodium, potassium or chloride could be clinically detrimental, and in patients presenting with:

- Hyperkalaemia.
 - Severe renal insufficiency (with oliguria/anuria).
 - Uncompensated cardiac or pulmonary failure.
 - Uncontrolled diabetes, other known glucose intolerances, hyperosmolar coma, hyperglycaemia and hyperlactataemia.
 - Hypersensitivity to maize products.
 - Disorders of fluid overload.
 - Hypersensitivity to the active ingredients or to any of the excipients listed in section 6.1.
- 4.4 Special warnings and precautions for use
- Fluid balance/renal function:**
Hyponatraemia
Depending on the tonicity of the solution, the volume and rate of infusion and on a patient's underlying clinical condition and capability to metabolise glucose, intravenous administration of glucose can cause electrolyte disturbances, most importantly hypo- or hyperosmotic hyponatraemia.
Treatment with intravenous fluids having a lower sodium concentration than the patient's serum sodium may cause hyponatraemia (see section 4.2).
Children, patients with reduced cerebral compliance, patients with non-osmotic vasopressin release (e.g. in acute illness, pain, trauma, post-operative stress, infections, burns and central nervous system diseases), patients with heart, liver and kidney diseases and patients exposed to vasopressin agonists and other medicines that can lower serum sodium (see section 4.5) are at particular risk of acute hyponatraemia.
Acute hyponatraemia can lead to acute hyponatraemic encephalopathy (brain oedema) characterised by headache, nausea, seizures, lethargy and vomiting. Patients with brain oedema are at particular risk of severe, irreversible and life-threatening brain injury.
Children, women of childbearing age and patients with reduced cerebral compliance (e.g. meningitis, intracranial bleeding, and cerebral contusion) are at particular risk of the severe and life-threatening brain swelling caused by acute hyponatraemia.

Risk of fluid and/or solute overload and electrolyte disturbances
The patient's clinical status and laboratory parameters (fluid balance, blood and urine electrolytes as well as acid-base balance) must be monitored during use of this solution.
Depending on the volume and rate of infusion, intravenous administration of MAINTENANCE SOLUTION FRESenius can cause fluid and/or solute overload resulting in overhydration/hypervolaemia, therefore high volume infusion must be specifically monitored in patients with cardiac, pulmonary or renal failure.

Use in patients with hypervolaemia or overhydration, or conditions that cause sodium retention and oedema
MAINTENANCE SOLUTION FRESenius should be used with great care in patients with hypertension, congestive heart failure, peripheral or pulmonary oedema, impaired renal function, severe renal insufficiency, pre-eclampsia, aldosteronism and in clinical states or other conditions in which there is sodium retention with oedema (also see section 4.5).
MAINTENANCE SOLUTION FRESenius should be used with care in patients with hypervolaemia, renal insufficiency, urinary tract obstruction, or impending or obvious cardiac decompensation.
Extraordinary electrolyte losses such as may occur during protracted nasogastric suction, vomiting, diarrhoea or gastrointestinal fistula drainage may necessitate additional electrolyte supplementation. Additional essential electrolytes, minerals and vitamins should be supplied as needed.

Care should be exercised in administering MAINTENANCE SOLUTION FRESenius to patients with renal or cardiovascular insufficiency, with or without congestive heart failure, particularly if they are post-operative or elderly.

Use in patients with severe renal impairment

In patients with diminished renal function, administration of MAINTENANCE SOLUTION FRESenius may result in sodium or potassium retention.

Electrolyte balance:

Use in patients with or at risk for hyperkalaemia
MAINTENANCE SOLUTION FRESenius should not be used in patients with hyperkalaemia, severe renal failure, and in conditions in which potassium retention is present.

Potassium therapy should be guided primarily by serial electrocardiograms, especially in patients receiving digoxin. Serum potassium levels are not necessarily indicative of tissue potassium levels. MAINTENANCE SOLUTION FRESenius should be used with caution in the presence of cardiac disease, particularly when accompanied by renal disease.

The plasma potassium level of the patient should be particularly closely monitored in patients at risk of hyperkalaemia. Solutions containing potassium salts should be administered with caution to patients with cardiac disease or conditions predisposing to hyperkalaemia such as renal or adrenocortical insufficiency, acute dehydration, or extensive tissue destruction as occurs with severe burns.

The following combinations are not recommended as they increase the concentration of potassium in the plasma and may lead to potentially fatal hyperkalaemia, notably in case of renal failure increasing the hyperkalaemic effects (see section 4.5):

- Angiotensin-converting enzyme (ACE) inhibitors and by extrapolation, angiotensin II receptor antagonists.
- Tacrolimus, ciclosporin.

Use in patients with hypocalcaemia

MAINTENANCE SOLUTION FRESenius contains no calcium, and an increase in plasma pH due to its alkalinising effect may lower the concentration of ionised (not protein-bound) calcium. MAINTENANCE SOLUTION FRESenius should be administered with particular caution to patients with hypocalcaemia.

Other warnings:

Hypersensitivity reactions
Hypersensitivity/infusion reactions, including anaphylactoid reactions, have been reported.
The infusion must be stopped immediately if any signs or symptoms of a suspected hypersensitivity reaction develop.
Appropriate therapeutic counter measures must be instituted as clinically indicated.
MAINTENANCE SOLUTION FRESenius should be used with caution in patients with known allergy to maize or maize products (see section 4.3).

Refeeding syndrome

For patients who are severely malnourished or have undergone a long period of starvation, caution should be exercised initially when administering dextrose. The dose may be gradually increased as dextrose metabolism improves. During long-term parenteral treatment, another convenient nutritive supply must be given to the patient.

Use in patients with or at risk for hyperglycaemia

MAINTENANCE SOLUTION FRESenius should be used with caution in patients with overt or known subclinical diabetes mellitus, or carbohydrate intolerance for any reason.

In diabetic patients, the amount of infused dextrose must be taken into account and insulin requirements may be modified. If hyperglycaemia occurs, rate of infusion should be adjusted, or insulin administered.

Due to the presence of dextrose, MAINTENANCE SOLUTION FRESenius is contraindicated in the first 24 hours following head trauma and blood glucose concentration should be closely monitored during intracranial hypertension episodes.
Hyperglycaemia has been implicated in increasing cerebral ischaemic brain damage and impairing recovery after acute ischaemic strokes. Caution is recommended in using dextrose-containing solutions in such patients.

Osmolarity

MAINTENANCE SOLUTION FRESenius for infusion is a hypertonic solution of electrolytes and dextrose (osmolarity: 382 mOsm/l). Administration of hypertonic solutions may cause venous irritation, including phlebitis. Hyperosmolar solutions should be administered with caution to patients with hyperosmolar states.

However, in the body, glucose-containing fluids can become extremely physiologically hypotonic due to rapid glucose metabolism (see section 4.2).

To minimise the risk of possible incompatibilities arising from mixing this solution with other additives that may be prescribed, the final infusate should be inspected for cloudiness or precipitation immediately after mixing, prior to administration, and periodically during administration.

If administration is controlled by a pumping device, care must be taken to discontinue pumping action before the container runs dry or air embolism may result.

MAINTENANCE SOLUTION FRESenius is intended for intravenous administration using sterile equipment.

For patients receiving a potassium supplement at greater than maintenance rates, frequent monitoring of serum potassium levels and serial ECGs are recommended.

If an adverse reaction does occur, discontinue the infusion, evaluate the patient, institute appropriate therapeutic countermeasures, and save the remainder of the fluid for examination if deemed necessary.

Use in paediatric patients

Safety and effectiveness of MAINTENANCE SOLUTION FRESenius in children have not been established.

The infusion rate and volume depend on the age, body mass, clinical and metabolic conditions of the patient and concomitant therapy, and should be determined by the consulting physician experienced in paediatric intravenous fluid therapy.

Newborns, especially those born premature and with low birth weight, are at increased risk of developing hypo- or hyperglycaemia and therefore need close monitoring during treatment with intravenous dextrose solutions to ensure adequate glycaemic control in order to avoid potential long-term adverse effects. Hypoglycaemia in the newborn can prolong seizures, coma and brain damage. Hyperglycaemia has been associated with intraventricular haemorrhage, late-onset bacterial and fungal infection, retinopathy of prematurity, necrotising enterocolitis, bronchopulmonary dysplasia, prolonged length of hospital stay and death.

Plasma electrolyte concentrations should be closely monitored in the paediatric population as this population may have impaired ability to regulate fluids and electrolytes.

The infusion of low sodium containing fluids together with the non-osmotic secretion of ADH may

PATIENT INFORMATION LEAFLET

SCHEDULING STATUS **S3**

MAINTENANCE SOLUTION FRESenius

Each 1 000 ml contains:	
Potassium chloride	1,938 g
Sodium chloride	1,5198 g
Dextrose anhydrous	50,0 g or
Dextrose monohydrate	55,0 g

Content of electrolytes per 1 000 ml:	
sodium ion (Na ⁺)	26 mmol/l
potassium ion (K ⁺)	26 mmol/l
chloride ion (Cl ⁻)	52 mmol/l

Contains sugar (dextrose).
Osmolarity: 382 mOsm/l
pH: 4,5
Energy: 840 kilojoules.

Read the entire leaflet carefully before you are given MAINTENANCE SOLUTION FRESenius

- Keep this leaflet. You may need to read it again.
- If you have further questions, please ask your doctor, pharmacist, nurse or other health care provider.

What is in this leaflet

1. What MAINTENANCE SOLUTION FRESenius is and what it is used for
2. What you need to know before you are given MAINTENANCE SOLUTION FRESenius
3. How you will be given MAINTENANCE SOLUTION FRESenius
4. Possible side effects
5. How to take MAINTENANCE SOLUTION FRESenius
6. Contents of the pack and other information.

1. What MAINTENANCE SOLUTION FRESenius is and what it is used for

MAINTENANCE SOLUTION FRESenius is a sterile intravenous solution providing fluid (water), energy in the form of sugar (dextrose) and electrolytes (sodium, potassium and chloride) to the body. Electrolytes are substances that become ions in solutions. Electrolytes are present in the human body. Maintenance of the correct electrolyte balance is essential for normal functioning of cells and organs.

MAINTENANCE SOLUTION FRESenius may be used intravenously to provide salts, energy and water to patients.

2. What you need to know before you are given MAINTENANCE SOLUTION FRESenius

You should not be given MAINTENANCE SOLUTION FRESenius:

- if you are hypersensitive (allergic) to potassium chloride, sodium chloride, dextrose or any of the other ingredients called corticosteroids that also occurs naturally in the body, which may be used as part of a medical test to check the functioning of your adrenal glands.
- if you are allergic to maize products, as dextrose may be manufactured from maize products
- if your body is retaining too much fluid
- if you have too much potassium in your blood
- if your kidneys are not functioning normally, and you fail to produce urine or have reduced urine production
- if you have heart and lung problems
- if you have diabetes, too much sugar or lactate in the blood.

Warnings and precautions

Tell your doctor or health care provider before being given the infusion:

- if you have low blood levels of sodium which can cause tiredness and confusion, muscle twitching, fits and coma (hyponatraemia)
- if you have swelling (oedema) and salt (sodium) retention
- if you have high levels of potassium in the blood (hyperkalaemia)
- if you have kidney failure and other kidney problems
- if you have liver problems
- if you have low levels of calcium in the blood (hypocalcaemia)
- if you have carbohydrate intolerance or diabetes
- if you are pregnant or suffer from pre-eclampsia, a pregnancy complication characterised by high blood pressure (see "Pregnancy and breastfeeding").

Too much MAINTENANCE SOLUTION FRESenius may cause congestion (swelling) of the lungs and limbs. Your doctor will examine you and do regular blood tests to decide how much and for how long you have to receive the treatment.

Other medicines and MAINTENANCE SOLUTION FRESenius

Always tell your health care provider if you are taking any other medicine. (This includes complementary or traditional medicines.)

Make sure that your doctor knows if you are also receiving the following:

- angiotensin used for high blood pressure (such as angiotensin-converting enzyme (ACE) inhibitors, angiotensin II receptor antagonists and telipressin)
- Medicines used to suppress the immune system (such as tacrolimus and ciclosporin).
- Medicines used for inflammation, asthma or arthritis (called corticosteroids, such as prednisolone).
- A hormone called corticotropin that also occurs naturally in the body, which may be used as part of a medical test to check the functioning of your adrenal glands.
- Medicines used for swelling, known as potassium-sparing diuretics such as amiloride, spironolactone, triamterene, alone or in combination).
- Medicines used for high blood sugar (such as chlorpropamide).
- Medicines used to treat epilepsy (such as carbamazepine).
- Medicines used to treat high levels of fat (cholesterol) in the blood (such as clofibrate).
- Cancer medicines (such as vincristine, ifosfamide and cyclophosphamide).
- Medicines used to treat depression (also known as selective serotonin reuptake inhibitors (SSRIs) and mental health problems.
- Medicines used to treat hyperactivity, also known as ADHD (such as methamphetamine).
- Medicines used for pain and inflammation, also known as NSAIDs.
- Medicines used for bed-wetting, called desmopressin.
- Medicines used during labour (oxytocin).
- Medicines used to manage anti-diuretic hormone deficiency and diabetes insipidus, known as vasopressin.

Ask your doctor, pharmacist or professional health care provider if you are not sure whether you are using or taking these medicines.

Pregnancy and breastfeeding

Safety in pregnancy and breastfeeding has not been established.

If you are pregnant or breastfeeding, think you may be pregnant or are planning to have a baby, please consult your doctor, pharmacist or other health care provider for advice before receiving MAINTENANCE SOLUTION FRESenius.

When receiving MAINTENANCE SOLUTION FRESenius during labour, especially if you are also receiving a medicine called oxytocin, there is a risk of reduced sodium levels in your blood.

If you received MAINTENANCE SOLUTION FRESenius while pregnant, your baby may be born with low blood glucose levels.

Driving and using machines

There is no information of the effects of MAINTENANCE SOLUTION FRESenius on the ability to drive and use machines. Ask your doctor if you are fit enough to drive or handle machines after release from hospital.

MAINTENANCE SOLUTION FRESenius contains dextrose

MAINTENANCE SOLUTION FRESenius contains 50 g dextrose anhydrous or 55 g dextrose monohydrate per 1 000 ml solution. This should be taken into account in patients with diabetes mellitus. Dextrose is a type of sugar, manufactured from maize. Tell your doctor if you are allergic to maize products.

3. How you will be given MAINTENANCE SOLUTION FRESenius

You will not be expected to administer MAINTENANCE SOLUTION FRESenius to yourself. Your doctor

result in hyponatraemia. Hyponatraemia can lead to headache, nausea, seizures, lethargy, coma, cerebral oedema and death. Therefore, acute symptomatic hyponatraemic encephalopathy is considered a medical emergency.

Use in elderly

In general, dose selection for an elderly patient should be cautious, usually starting at the low end of the dosing range, reflecting the greater frequency of decreased hepatic, renal, or cardiac function, and of concomitant disease or other medicinal therapy.

These medicinal therapies are known to be substantially excreted by the kidney, and the risk of toxic reactions to these medicines may be greater in patients with impaired renal function. Because elderly patients are more likely to have decreased renal function, care should be taken in dose selection, and it may be useful to monitor renal function. See section 4.4.

4.5 Interaction with other medicines and other forms of interaction

Interaction with sodium:

- MAINTENANCE SOLUTION FRESenius should be administered with caution to patients receiving corticosteroids or corticotropin, or to patients with salt retaining disorders.

Interaction with potassium:

- Potassium-sparing diuretics (amiloride, spironolactone, triamterene, alone or in combination).
- Angiotensin-converting enzyme (ACE) inhibitors and angiotensin II receptor antagonists.
- Tacrolimus and ciclosporin increase the concentration of potassium in the plasma and the combination may lead to potentially fatal hyperkalaemia, notably in case of renal failure contributing to the hyperkalaemia.

Medicines leading to an increased vasopressin effect:

The medicines listed below increase the vasopressin effect, leading to reduced renal electrolyte free water excretion and increase the risk of hospital-acquired hyponatraemia following inappropriately balanced treatment with IV fluids (see sections 4.2, 4.4 and 4.8).

- Medicines stimulating vasopressin release, such as chlorpropamide, clofibrate, carbamazepine, vincristine, selective serotonin reuptake inhibitors, 3,4-methylenedioxy-*N*-methamphetamine, ifosfamide, antipsychotics, narcotics.
- Medicines potentiating vasopressin action such as chlorpropamide, NSAIDs and cyclophosphamide.
- Vasopressin analogues such as desmopressin, oxytocin, vasopressin and telipressin.

Other medicines increasing the risk of hyponatraemia also include diuretics in general and antiepileptics such as oxcarbazepine.

Medicines leading to an increased vasopressin effect:

The medicines listed below increase the vasopressin effect, leading to reduced renal electrolyte free water excretion and increase the risk of hospital-acquired hyponatraemia following inappropriately balanced treatment with IV fluids (see sections 4.2, 4.4 and 4.8).

- Medicines stimulating vasopressin release, such as chlorpropamide, clofibrate, carbamazepine, vincristine, selective serotonin reuptake inhibitors, 3,4-methylenedioxy-*N*-methamphetamine, ifosfamide, antipsychotics, narcotics.
- Medicines potentiating vasopressin action such as chlorpropamide, NSAIDs and cyclophosphamide.
- Vasopressin analogues such as desmopressin, oxytocin, vasopressin and telipressin.

Other medicines increasing the risk of hyponatraemia also include diuretics in general and antiepileptics such as oxcarbazepine.

4.6 Fertility, pregnancy and lactation

Safety during pregnancy and lactation has not been established.

When MAINTENANCE SOLUTION FRESenius is administered to pregnant women during labour, particularly if administered in combination with oxytocin, there may be an increased risk for hyponatraemia (see sections 4.4, 4.5 and 4.8).

Intrapartum maternal intravenous infusion of MAINTENANCE SOLUTION FRESenius may result in fetal insulin production, with an associated risk of fetal hyperglycaemia and metabolic acidosis as well as rebound hypoglycaemia in the neonate.

4.7 Effects on ability to drive and use machines

There is no information of the effects of MAINTENANCE SOLUTION FRESenius on the ability to drive and use machines. Patients should be advised to take special care before performing tasks requiring their attention, until they know how MAINTENANCE SOLUTION FRESenius will affect them.

4.8 Undesirable effects

Immune system disorders:

Frequency not known:
Hypersensitivity/infusion reaction including anaphylactoid reaction.

Metabolism and nutrition disorders:

Frequency not known:
Hyperkalaemia, hyperglycaemia, hypervolaemia, hospital-acquired hyponatraemia, electrolyte disturbance.

Nervous system disorders:

Frequency not known:
Mental confusion, seizures, hyponatraemic encephalopathy.

Cardiac disorders:

Frequency not known:
Hypotension, cardiac dysrhythmias, heart block, electrocardiographic abnormalities and cardiac arrest.

Respiratory, thoracic and mediastinal disorders:

Frequency not known:
Respiratory paralysis.

Gastrointestinal disorders:

Frequency not known:
Nausea, vomiting, abdominal pain and diarrhoea, intestinal ileus and dilatation.

Musculoskeletal and connective tissue disorders:

Frequency not known:
Paraesthesia of the extremities, areflexia, respiratory or muscular paralysis, disruption of neuromuscular function and weakness.

General disorders and administration site conditions:

Frequency not known:
Infusion site adverse reactions which may occur because of MAINTENANCE SOLUTION FRESenius or the technique of administration include a burning sensation, a febrile response (pyrexia), infection, irritation and pain at the site of injection, venous thrombosis or thrombophlebitis extending from the site of injection, extravasation, and hypervolaemia.

Too rapid rate of infusion of MAINTENANCE SOLUTION FRESenius may cause local pain and venous irritation. The rate of administration should be adjusted according to tolerance. Use of the largest peripheral vein and a small bore catheter is recommended. See section 4.2. Symptoms may result from an excess or deficit of one or more of the ions present in the solution; therefore, frequent monitoring of electrolyte levels is essential.

Hypermnatraemia may be associated with oedema and exacerbation of congestive heart failure due to the retention of water, resulting in an expanded extracellular fluid volume.

If MAINTENANCE SOLUTION FRESenius is infused in large amounts, chloride ions may cause a loss of bicarbonate ions, resulting in an acidifying effect.

Reporting of suspected adverse reactions:

Health care providers are asked to report any suspected adverse drug reactions to the Holder of the Certificate of Registration at the following email address: safety.fksa@fresenius-kabi.com and to the relevant medicine's regulatory authority in the country where the product is marketed.

Reporting suspected adverse reactions after authorisation of MAINTENANCE SOLUTION FRESenius is important. It allows continued monitoring of the benefit/risk balance of MAINTENANCE SOLUTION FRESenius. Health care providers are asked to report any suspected adverse reactions to SAHPRA via the "6.04 Adverse Drug Reactions Reporting Form", found online under SAHPRA's publications: <https://www.sahpra.org.za/Publications/Index/8>.

4.9 Overdose

In the event of a fluid or solute overload during parenteral therapy, re-evaluate the patient's condition and institute appropriate corrective treatment.

Too fast an infusion rate may lead to water and sodium overload with a risk of oedema, particularly when there is a defective renal sodium excretion. In this case renal dialysis may be necessary.

Excessive administration, rapid infusion or prolonged administration of glucose may lead to hyperglycaemia.

In the event of overdose with potassium-containing solutions, discontinue the infusion immediately and institute corrective therapy to reduce serum potassium levels.

Excessive administration of potassium may lead to the development of hyperkalaemia, especially in patients with renal impairment. Symptoms include paraesthesia of the extremities, muscle weakness, paralysis, cardiac dysrhythmias, heart block, cardiac arrest, and mental confusion. Treatment of hyperkalaemia involves the administration of calcium, insulin (with glucose), sodium bicarbonate, exchange resins or dialysis.

Excessive administration of chloride salts may cause a loss of bicarbonate with an acidifying effect.

Acute treatment:

The infusion should be interrupted immediately. Administration of diuretics and continuous monitoring of serum electrolytes, correction of electrolyte balance and acid-base balance.

When assessing an overdose, any additives in the solution must also be considered.

The effects of an overdose may require immediate medical attention and treatment.

5. PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

Category and class: A 24 Mineral substitutes, electrolytes.
Pharmacotheapeutic group: Electrolytes with carbohydrates.

ATC code: B05BB02.

MAINTENANCE SOLUTION FRESenius provides water, electrolytes and energy.

5.2 Pharmacokinetic properties

After intravenous administration dextrose exhibits fast (approximately 20 minutes) and slow phases of equilibrium. Distribution is largely through extracellular water, and intracellular water of the liver.

In the post-operative state endogenous glucose production equals the amount of glucose taken up and metabolised by all the tissues and is about $2,3 \pm 0,1 \text{ mg/kg}^{-1} \text{ min}^{-1}$.

Sodium homeostasis is complex and closely associated with fluid balance. Osmolality and volume of extracellular fluid are tightly regulated. Small changes in osmolality (plasma sodium concentrations) are corrected by alterations of extracellular volume. The balance of plasma osmolality is achieved by the secretion or suppression of antidiuretic hormone (ADH; vasopressin), which primarily controls water excretion by the kidney.

A normal concentration of potassium in plasma is 3,5 to 5,0 mmol per litre, but factors influencing the intracellular and extracellular shifts, such as acid-base disturbances, can distort the relationship between plasma concentrations and total body stores.

6. PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Water for injection

6.2 Incompatibilities

Before adding a medicine to MAINTENANCE SOLUTION FRESenius, verify that it is soluble and stable in water at the pH of MAINTENANCE SOLUTION FRESenius (pH 4,5).

MAINTENANCE SOLUTION FRESenius should not be administered concomitantly with blood in the same infusion set due to the risk of clotting of red blood cells.

6.3 Shelf life

24 months.

6.4 Special precautions for storage

</

